



AZSTARYS® is the first
and only ADHD
treatment of its kind

**Helps control ADHD symptoms
from the morning until evening.**



Actor portrayal.

What is AZSTARYS?

AZSTARYS is a central nervous system (CNS) stimulant prescription medicine for the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in people 6 years of age and older. AZSTARYS may help increase attention and decrease impulsiveness and hyperactivity in people 6 years of age and older with ADHD.

It is not known if AZSTARYS is safe and effective in children younger than 6 years of age.

IMPORTANT SAFETY INFORMATION

Abuse, misuse, and addiction. AZSTARYS has a high chance for abuse and misuse and may lead to substance use problems, including addiction. Misuse and abuse of AZSTARYS, other methylphenidate containing medicines, and amphetamine containing medicines, can lead to overdose and death. The risk of overdose and death is increased with higher doses of AZSTARYS or when it is used in ways that are not approved, such as snorting or injection.

- Your healthcare provider (HCP) should check you or your child's risk for abuse, misuse, and addiction before starting treatment with AZSTARYS. Tell your HCP if you or your child have ever abused or been dependent on alcohol, prescription medicines, or street drugs.
- AZSTARYS may lead to physical dependence after prolonged use, even if taken as directed by your HCP.
- Do not give AZSTARYS to anyone else because it may cause death or harm them. Keep AZSTARYS in a safe place to protect it from theft and properly dispose of any unused medicine.

AZSTARYS is a federally controlled substance (CII) because it contains dextmethylphenidate that can be a target for people who abuse prescription medicines or street drugs.

Please see additional Important Safety Information throughout and Full Prescribing Information and Medication Guide.



**serdexmethylphenidate
and dextmethylphenidate**

26.1/5.2mg • 39.2/7.8mg • 52.3/10.4mg capsules

Are your ADHD symptoms under control?

A day with ADHD symptoms can be filled with challenges. It's important to discuss with your doctor how these symptoms affect you at work, at home, or in social situations.

I forget appointments and lose important things.

It's hard to stay focused in conversations and activities.

I lose track of time and end up running late in the morning.

I can't control the feeling of restlessness.

I talk a lot and may interrupt others.

Keeping organized and productive at work is a struggle.



Actor portrayal.

★ ★ **Talk with your doctor about your ADHD symptoms and your options to help keep them in check.**

This information is not meant to diagnose ADHD.

IMPORTANT SAFETY INFORMATION (continued)

Who should not take AZSTARYS?

Do not take AZSTARYS if you or your child are:

- allergic to serdexmethylphenidate, methylphenidate, or any of the ingredients in AZSTARYS.
- taking or have stopped taking within the past 14 days a medicine used to treat depression called a monoamine oxidase inhibitor (MAOI).

Adult ADHD by the numbers

4.4% of adults in the United States are estimated to **have ADHD**.

That's about **10.5 million** adults.

60% of people with ADHD are estimated to **carry it into adulthood**.

41% to 55% of families with at least 1 child with ADHD **also have at least 1 parent with ADHD**.

IMPORTANT SAFETY INFORMATION (continued)

AZSTARYS may cause serious side effects, including:

- **Risks for people with serious cardiac disease.** Sudden death has happened in people who have heart defects or other serious heart disease. Your HCP should check you or your child carefully for heart problems before starting treatment with AZSTARYS. Tell your HCP if you or your child have any heart problems, heart disease, or heart defects. **Call your HCP or go to the nearest hospital emergency room right away if you or your child have any signs of heart problems such as chest pain, shortness of breath, or fainting during treatment with AZSTARYS.**
- **Increased blood pressure and heart rate.** Your HCP should check you or your child's blood pressure and heart rate regularly during treatment with AZSTARYS.

Please see additional Important Safety Information throughout and Full Prescribing Information and Medication Guide.


azstarlys®
serdexmethylphenidate
and dexamethylphenidate
26.1/5.2mg • 39.2/7.8mg • 52.3/10.4mg capsules

Discover what makes **AZSTARYS** different

AZSTARYS is pronounced *az-star'-is*.

AZSTARYS has 2 forms of medicine in 1 capsule

Dexmethylphenidate

(deks-meth-uhl-fen-uh-date), or **d-MPH**

An immediate-release
medicine that kicks in quickly

+

Serdexmethylphenidate

(sur-deks-meth-uhl-fen-uh-date), or **SDX**

A different way to provide continuous
medicine throughout the day

★ ★
Talk with your doctor to see if
AZSTARYS may be right for you.

Visit [AZSTARYS.com](https://www.azstarys.com) to learn more.




IMPORTANT SAFETY INFORMATION (continued)

AZSTARYS may cause serious side effects, including: (continued)

- **Mental (psychiatric) problems, including:** new or worse behavior and thought problems, new or worse bipolar illness, new psychotic symptoms (such as hearing voices, or seeing or believing things that are not real), or new manic symptoms. Tell your HCP about any mental problems you or your child have, or about a family history of suicide, bipolar illness, or depression. **Call your HCP right away if you or your child have any new or worsening mental symptoms or problems during treatment with AZSTARYS, especially hearing voices, seeing or believing things that are not real, or new manic symptoms.**

AZSTARYS is designed
to help in 3 ways, from
morning to evening



1 Starts working quickly to
help in the morning

2 Helps control ADHD
symptoms throughout
the day

3 Wears off gradually in
the evening

IMPORTANT SAFETY INFORMATION (continued)

AZSTARYS may cause serious side effects, including: (continued)

- **Painful and prolonged erections (priapism).** Priapism has happened in males who take products that contain methylphenidate. If you or your child develop priapism, get medical help right away.
- **Circulation problems in fingers and toes (peripheral vasculopathy, including Raynaud's phenomenon).** Tell your HCP if you or your child have numbness, pain, skin color change, or sensitivity to temperature in the fingers or toes. Call your HCP right away if you or your child have any signs of unexplained wounds appearing on fingers or toes during treatment with AZSTARYS.

Please see additional Important
Safety Information throughout
and Full Prescribing Information
and Medication Guide.


serdexmethylphenidate
and dexmethylphenidate
26.1/5.2mg • 39.2/7.8mg • 52.3/10.4mg capsules

How to take AZSTARYS

Take AZSTARYS exactly as prescribed

There are different options for taking AZSTARYS as part of the morning routine



Swallow whole

An AZSTARYS capsule can be swallowed whole with water or other liquids.

or



Sprinkle into water or onto applesauce

The capsule can be opened and its contents can be sprinkled into 2 ounces (50 mL) of water or onto 2 tablespoons of applesauce.

All of the water or applesauce mixture should be swallowed within 10 minutes. **Do not** store the water or applesauce mixture.

★ ★ AZSTARYS can be taken
with or without food.



Remembering to take AZSTARYS

AZSTARYS should be taken in the morning. It may be helpful to set an alert to remind you to take your AZSTARYS each day.

IMPORTANT SAFETY INFORMATION (continued)

AZSTARYS may cause serious side effects, including: (continued)

- **Slowing of growth (height and weight) in children.** Children should have their height and weight checked often during treatment with AZSTARYS. AZSTARYS treatment may be stopped if your child is not growing or gaining weight.
- **Eye problems (increased pressure in the eye and glaucoma).** Tell your HCP if you or your child have eye problems, including increased pressure in your eye, glaucoma, or problems with your close-up vision (farsightedness). Call your HCP right away if you or your child develop changes in your vision, eye pain, swelling, or redness.

AZSTARYS comes in 3 dosage strengths

- Each dosage strength is a different color
- All the capsules are the same size



Capsules are not true to size.

IMPORTANT SAFETY INFORMATION (continued)

AZSTARYS may cause serious side effects, including: (continued)

- **New or worsening tics or worsening Tourette's syndrome.** Tell your HCP if you or your child get any new or worsening tics or worsening Tourette's syndrome during treatment with AZSTARYS.

Before taking AZSTARYS, tell your HCP about all medical conditions including if you or your child are pregnant or plan to become pregnant, because it is not known if AZSTARYS will harm the unborn baby, or are breastfeeding or plan to breastfeed, as AZSTARYS passes into breast milk.

Tell your HCP about all the medicines you or your child take, especially if you or your child take blood pressure medicines (anti-hypertensive). Do not start any new medicine during treatment with AZSTARYS without talking to your HCP first.

Please see additional Important Safety Information throughout and Full Prescribing Information and Medication Guide.

azstarys [®] _{II}
serdexmethylphenidate
and dexamethylphenidate
26.1/5.2mg • 39.2/7.8mg • 52.3/10.4mg capsules



Actor portrayals.



★ Visit **AZSTARYS.com** to learn about copay assistance.

IMPORTANT SAFETY INFORMATION (continued)

What are the most common side effects of AZSTARYS?

The most common side effects of AZSTARYS include:

- decreased appetite
- nausea
- indigestion
- weight loss
- dizziness
- mood swings
- increased blood pressure
- trouble sleeping
- vomiting
- stomach pain
- anxiety
- irritability
- increased heart rate

These are not all the possible side effects of AZSTARYS. Call your HCP for medical advice about side effects.

Online ADHD resources

Connecting with others who have been in your shoes and getting more information about ADHD can be valuable in helping on your ADHD journey. These websites have plenty of useful information about ADHD in adults.

Attention Deficit Disorder Association (ADDA)

<https://www.add.org>

Children and Adults with Attention-Deficit/Hyperactivity Disorder (CHADD)

<https://www.chadd.org>

ADDitude

<https://www.additudemag.com>

National Institute of Mental Health

<https://www.nimh.nih.gov/health/topics/attention-deficit-hyperactivity-disorder-adhd>

American Professional Society of ADHD and Related Disorders

<https://www.apsard.org>

IMPORTANT SAFETY INFORMATION (continued)

What is AZSTARYS?

AZSTARYS is a central nervous system (CNS) stimulant prescription medicine for the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in people 6 years of age and older. AZSTARYS may help increase attention and decrease impulsiveness and hyperactivity in people 6 years of age and older with ADHD.

It is not known if AZSTARYS is safe and effective in children younger than 6 years of age.

Please see additional Important Safety Information throughout and Full Prescribing Information and Medication Guide.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.



serdexmethylphenidate
and dexamethylphenidate

26.1/5.2mg • 39.2/7.8mg • 52.3/10.4mg capsules

IMPORTANT SAFETY INFORMATION

Abuse, misuse, and addiction. AZSTARYS has a high chance for abuse and misuse and may lead to substance use problems, including addiction. Misuse and abuse of AZSTARYS, other methylphenidate containing medicines, and amphetamine containing medicines, can lead to overdose and death. The risk of overdose and death is increased with higher doses of AZSTARYS or when it is used in ways that are not approved, such as snorting or injection.

- Your healthcare provider (HCP) should check you or your child's risk for abuse, misuse, and addiction before starting treatment with AZSTARYS. Tell your HCP if you or your child have ever abused or been dependent on alcohol, prescription medicines, or street drugs.
- AZSTARYS may lead to physical dependence after prolonged use, even if taken as directed by your HCP.
- Do not give AZSTARYS to anyone else because it may cause death or harm them. Keep AZSTARYS in a safe place to protect it from theft and properly dispose of any unused medicine.

AZSTARYS is a federally controlled substance (CII) because it contains dexmethylphenidate that can be a target for people who abuse prescription medicines or street drugs.

Who should not take AZSTARYS?

Do not take AZSTARYS if you or your child are:

- allergic to serdexmethylphenidate, methylphenidate, or any of the ingredients in AZSTARYS.
- taking or have stopped taking within the past 14 days a medicine used to treat depression called a monoamine oxidase inhibitor (MAOI).

AZSTARYS may cause serious side effects, including:

- **Risks for people with serious cardiac disease.** Sudden death has happened in people who have heart defects or other serious heart disease. Your HCP should check you or your child carefully for heart problems before starting treatment with AZSTARYS. Tell your HCP if you or your child have any heart problems, heart disease, or heart defects. **Call your HCP or go to the nearest hospital emergency room right away if you or your child have any signs of heart problems such as chest pain, shortness of breath, or fainting during treatment with AZSTARYS.**
- **Increased blood pressure and heart rate.** Your HCP should check you or your child's blood pressure and heart rate regularly during treatment with AZSTARYS.
- **Mental (psychiatric) problems, including:** new or worse behavior and thought problems, new or worse bipolar illness, new psychotic symptoms (such as hearing voices, or seeing or believing things that are not real), or new manic symptoms. Tell your HCP about any mental problems you or your child have, or about a family history of suicide, bipolar illness, or depression. **Call your HCP right away if you or your child have any new or worsening mental symptoms or problems during treatment with AZSTARYS, especially hearing voices, seeing or believing things that are not real, or new manic symptoms.**

IMPORTANT SAFETY INFORMATION (continued)

AZSTARYS may cause serious side effects, including: (continued)

- **Painful and prolonged erections (priapism).** Priapism has happened in males who take products that contain methylphenidate. If you or your child develop priapism, get medical help right away.
- **Circulation problems in fingers and toes (peripheral vasculopathy, including Raynaud's phenomenon).** Tell your HCP if you or your child have numbness, pain, skin color change, or sensitivity to temperature in the fingers or toes. Call your HCP right away if you or your child have any signs of unexplained wounds appearing on fingers or toes during treatment with AZSTARYS.
- **Slowing of growth (height and weight) in children.** Children should have their height and weight checked often during treatment with AZSTARYS. AZSTARYS treatment may be stopped if your child is not growing or gaining weight.
- **Eye problems (increased pressure in the eye and glaucoma).** Tell your HCP if you or your child have eye problems, including increased pressure in your eye, glaucoma, or problems with your close-up vision (farsightedness). Call your HCP right away if you or your child develop changes in your vision, eye pain, swelling, or redness.
- **New or worsening tics or worsening Tourette's syndrome.** Tell your HCP if you or your child get any new or worsening tics or worsening Tourette's syndrome during treatment with AZSTARYS.

Before taking AZSTARYS, tell your HCP about all medical conditions including if you or your child are pregnant or plan to become pregnant, because it is not known if AZSTARYS will harm the unborn baby, or are breastfeeding or plan to breastfeed, as AZSTARYS passes into breast milk.

Tell your HCP about all the medicines you or your child take, especially if you or your child take blood pressure medicines (anti-hypertensive). Do not start any new medicine during treatment with AZSTARYS without talking to your HCP first.

What are the most common side effects of AZSTARYS?

The most common side effects of AZSTARYS include:

- | | |
|----------------------------|------------------------|
| • decreased appetite | • trouble sleeping |
| • nausea | • vomiting |
| • indigestion | • stomach pain |
| • weight loss | • anxiety |
| • dizziness | • irritability |
| • mood swings | • increased heart rate |
| • increased blood pressure | |

These are not all the possible side effects of AZSTARYS. Call your HCP for medical advice about side effects.

Please see additional Important Safety Information throughout and Full Prescribing Information and Medication Guide.


serdexmethylphenidate
and dexmethylphenidate
26.1/5.2mg • 39.2/7.8mg • 52.3/10.4mg capsules

AZSTARYS is the
first and only ADHD
treatment of its kind



**Copay savings and support are
available for eligible patients.***

Visit **AZSTARYS.com** or call **1-800-910-8432**
to learn about copay assistance.



Scan the QR code using your phone's
camera to see savings information.

*Eligibility required. See Terms and Conditions at [AZSTARYS.com](https://www.azstarys.com).

IMPORTANT SAFETY INFORMATION (continued)

What is AZSTARYS?

AZSTARYS is a central nervous system (CNS) stimulant prescription medicine for the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in people 6 years of age and older. AZSTARYS may help increase attention and decrease impulsiveness and hyperactivity in people 6 years of age and older with ADHD.

It is not known if AZSTARYS is safe and effective in children younger than 6 years of age.

Please see additional Important Safety Information throughout and Full Prescribing Information and Medication Guide.

**You are encouraged to report
negative side effects of prescription
drugs to the FDA. Visit
www.fda.gov/medwatch, or call
1-800-FDA-1088.**

Corium

© 2023 Corium, LLC. All Rights Reserved.
AZS-0326 (v2.0) 11/23


azstarys®

**serdexmethylphenidate
and dexamethylphenidate**

26.1/5.2mg • 39.2/7.8mg • 52.3/10.4mg capsules